

Fifteen-minute consultation: Approach to the adolescent presenting with hirsutism

Alexandra Selina Cross ¹, Manar Moustafa,² Charlotte Jane Elder ^{1,3}

¹Paediatric Endocrinology, Sheffield Children's Hospital NHS Foundation Trust, Sheffield, UK

²Paediatric Dermatology, Sheffield Children's NHS Foundation Trust, Sheffield, UK

³Department of Oncology and Metabolism, The University of Sheffield, Sheffield, UK

Correspondence to

Dr Charlotte Jane Elder, Academic Unit of Child Health, The University of Sheffield, Sheffield, S10 2TN, UK; C.J.Elder@sheffield.ac.uk

Received 24 August 2022

Accepted 1 January 2023

Published Online First

19 January 2023

ABSTRACT

Hirsutism, unwanted terminal hair growth in androgen-dependent areas, is a common presentation to general paediatricians, dermatologists and endocrinologists. Polycystic ovarian syndrome is the most common cause but can be challenging to diagnose in young people due to the significant overlap of features with the healthy adolescent population.

There are other rare, but important, causes to consider such as non-classic congenital adrenal hyperplasia and androgen-secreting tumours. Hirsutism carries a significant psychological burden for those living with it. This 15 min consultation piece describes the causes of hirsutism, introduces a novel assessment tool and suggests an approach to investigations and management, including signposting to psychological support.

INTRODUCTION

Hirsutism is the growth of excessive terminal hair in women in an adult male distribution.¹ Terminal hair is thicker and darker than small, fair vellus hair. Growth of terminal hair, from vellus hair, is androgen dependent and influenced by plasma androgen levels and hair follicle androgen sensitivity.¹ Hirsutism should be differentiated from hypertrichosis, generalised excess hair growth in non-androgen-dependent areas, for example, the forearms.² Hirsutism is a common condition, estimated to affect 3%–20% of adolescent girls³ with associated psychosocial morbidity.¹

WHAT CAUSES HIRSUTISM?

Idiopathic hirsutism, with or without hyperandrogenism, is common, but the most common cause in adolescents is polycystic ovarian syndrome (PCOS).^{3 4} The diagnosis of PCOS is challenging in adolescents due to a considerable overlap with the healthy population as features

like acne and irregular menstrual cycles are commonplace. Modified diagnostic criteria should be used.⁵ Rarer conditions, including non-classic congenital adrenal hyperplasia (NCCAH), androgen-secreting tumours and other endocrinopathies may present with hirsutism and are important to consider (table 1).³

ASSESSMENT

Adolescents presenting with excess hair growth require a thorough evaluation. We have developed a clinical assessment proforma for this purpose (figure 1).

History

'Patient important hirsutism' is any excessive terminal hair growth causing the patient sufficient distress to seek treatment. Consultations should be conducted sensitively, with the clinician proactively assessing the psychosocial impact of the unwanted hair, including the patient's perception of their hair growth and effects on their confidence, self-esteem, relationships, school and activities. It is important to explore the patient's (and parent's/carer's) expectations to aid empathic discussion of realistic and achievable treatment outcomes. Detailed history should be sought about the age/stage of puberty at onset and speed of hair growth (prepubertal onset and rapid progression are concerning for androgen secreting tumours).³ Often patients have removed the hair, so a description of the hair and its distribution will help assess severity and differentiate it from hypertrichosis. The efficacy and side effects of hair removal techniques should be noted. Ask appropriate questions to identify features of the differential diagnoses in table 1.

Examination

Hirsutism is typically assessed using the modified Ferriman-Gallwey (mFG) score (figure 2), with hair growth recorded in



© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Cross AS, Moustafa M, Elder CJ. *Arch Dis Child Educ Pract Ed* 2024;**109**:66–72.

Table 1 Causes of hirsutism in adolescents with the key diagnostic features and supportive investigations

Aetiology	Relative prevalence in patients with hirsutism	Diagnostic features	Supportive investigations
Polycystic ovary syndrome	20%–40% in adolescent girls ³	<ul style="list-style-type: none"> ▶ Clinical (hirsutism, severe cystic acne) and/or biochemical evidence of hyperandrogenism⁴ and irregular menses*. ▶ Associated but non-diagnostic features include obesity and insulin resistance (acanthosis nigricans).⁴ ▶ Ovarian USS should not be routinely performed until 8 years post menarche as the polycystic ovarian morphology of large, multifollicular ovaries are common in adolescents.^{7,14} 	<ul style="list-style-type: none"> ▶ ↑ testosterone (although may be normal) or ↑ FAI (FAI calculation=(total testosterone/SHBG)×100). ▶ Diagnosis of exclusion. ▶ Patients with amenorrhoea/ oligomenorrhoea for less than 3 years,⁷ who have evidence of hyperandrogenism are 'at risk' of PCOS and should be followed up.⁴
Idiopathic hyperandrogenism	10%–15% ¹³	<ul style="list-style-type: none"> ▶ No other identifiable cause of elevated androgen levels. ▶ Normal menses. 	<ul style="list-style-type: none"> ▶ ↑ testosterone or ↑ FAI. ▶ Diagnosis of exclusion.
Idiopathic hirsutism	6%–10% ^{13,15} (half of mild hirsutism) ⁸	<ul style="list-style-type: none"> ▶ Normal androgen levels and menses. ▶ No features of other endocrine disorder. ▶ Often familial or related to ethnicity.¹¹ 	<ul style="list-style-type: none"> ▶ Normal testosterone. ▶ Diagnosis of exclusion.
NCCAH	4%–14% ^{3,15}	<ul style="list-style-type: none"> ▶ Family history of CAH, ↑ risk ethnicity (Ashkenazi Jew, Hispanic, Slav). ▶ Premature pubarche (before age of 8 years). ▶ Accelerated growth. ▶ Signs of virilisation (clitoromegaly, increased muscle mass, voice change). 	<ul style="list-style-type: none"> ▶ Indicative USP. ▶ ↑ 17OHP, early morning in follicular phase. ▶ 17OHP level >30 nmol/L basal or after Synacthen stimulation diagnostic.⁹ ▶ 17OHP levels >6 nmol/L in prepuberty.⁹
Androgen-secreting tumours	0.2% (half of which are malignant) ³	<ul style="list-style-type: none"> ▶ Evidence of virilisation. ▶ Rapid onset. ▶ Abdominal/pelvic mass. ▶ Progressive hair growth despite treatment.³ 	<ul style="list-style-type: none"> ▶ Raised testosterone (>5 nmol/L or >2–3×ULNR). ▶ Indicative USP.³ ▶ ↑ DHEAS (>19 mmol/L) (adrenal source). ▶ ↑ androstenedione (ovarian source). ▶ Adrenal or ovarian mass on USS or MRI.³
Other		<ul style="list-style-type: none"> ▶ Other symptoms likely to predominate. ▶ Cushing syndrome: lack of height gain despite ↑ weight, central obesity, plethoric facies, proximal muscle weakness. ▶ Pregnancy: amenorrhoea, nausea and fatigue. ▶ Hyperprolactinaemia: galactorrhoea, amenorrhoea, features of space-occupying lesion, for example, bitemporal hemianopia. 	<ul style="list-style-type: none"> ▶ Urinary cortisol, late night salivary cortisol, dexamethasone suppression test.¹⁶ ▶ ↑ urine or plasma beta HCG. ▶ ↑ prolactin (exclude macroprolactin).¹⁷
Drugs		Access to drugs including <ul style="list-style-type: none"> ▶ Anabolic steroids, testosterone and valproate.⁸ 	<ul style="list-style-type: none"> ▶ USP.

*Irregular menses are normal in the first year after menarche, although cycles <19 days or >90 days are considered abnormal.²

CAH, congenital adrenal hyperplasia; DHEAS, dehydroepiandrosterone sulfate; FAI, Free Androgen Index; HCG, human chorionic gonadotropin; NCCAH, non-classic congenital adrenal hyperplasia; 17OHP, 17-hydroxyprogesterone; PCOS, polycystic ovary syndrome; SHBG, sex hormone-binding globulin; ULNR, upper limit normal range; USP, urinary steroid profile; USS, ultrasound scan.

nine androgen-sensitive areas: upper lip, chin, chest, upper arms, upper back, lower back, upper abdomen, lower abdomen and thighs. Each area is scored from 0 to 4, with 0 denoting absence of terminal hair, 1 representing minimal hirsutism and 4 signifying frank virilisation, equivalent to a 'hairy' man. A total score of 8 is suggested as the threshold for mild hirsutism, 16–24 for moderate and >24 for severe hirsutism.⁶ The mFG score has not been validated in adolescents, who may have lower scores due to reduced exposure time to androgens.⁷ No standardised cut-offs for different ethnicities have been agreed. It is suggested higher thresholds are used in patients of Mediterranean or South Asian origin and lower scores in those of East Asian heritage (figure 1)^{1,6,8}; however, specific cut-offs have not been validated in adolescents from different ethnicities.³ The mFG does not account for patients with extensive but localised hair growth. Each body area is equally represented, meaning patients affected

by only facial hair growth may not score highly but are likely to experience distress and would benefit from treatment.¹

Investigations

Investigation of patients presenting with hirsutism should be guided by the clinician's assessment, mFG score, and presence of red flag features of hyperandrogenism or an endocrinopathy (figure 3). In patients with isolated hair growth or low mFG scores, biochemical investigation is usually not required. Most patients will have mild hirsutism and, in the absence of red flag features, require only testosterone and sex hormone-binding globulin (SHBG) enabling calculation of the Free Androgen Index (FAI), a surrogate marker of free and therefore active testosterone. There is no universally agreed normal range for FAI, so local laboratory cut-offs need to be employed. 17-Hydroxyprogesterone (17OHP) should be measured in patients with raised

Hirsutism Assessment Proforma

Name:
 DOB:
 Hospital number:
 (Or use patient label)

History

Hair growth

Anatomical location of hair
 Age & pubertal stage at onset
 Speed of onset and progression
 Hair removal techniques tried, effect and side effect
 Time since last hair removal

Psychosocial

Effect on patient QoL and mental health
 (e.g., bullying, self-esteem, social withdrawal, self-harm, support system)

Menstrual history

Age of menarche
 Cycle duration/ frequency
 Regular or irregular*? Regular Irregular
*Irregular = 1-3y post menarche: cycle length <21d or >45d. >3y post menarche: <21d or >35d or <8 cycles/ year)

Examination

BP: Weight: Height: BMI:
 Centile: Centile: Centile: Centile:

Tanner puberty stage

Document if consent given, chaperone offered, accepted and name
 Breast Pubic hair Axillary hair
 Signs of virilisation? Yes No

Symptoms

Features of hyperandrogenism? Yes No
 (E.g., deepening of voice, increased muscle bulk, shrinkage of breasts, severe treatment resistant acne, clitoromegaly, androgenic alopecia)
 Features of endocrinopathies? Yes No
 (E.g. galactorrhoea, features of Cushing's syndrome or thyroid dysfunction)
 Other symptoms? Yes No
 (E.g., headache, visual change, nausea, weight change, excessive exercise, significant stress, change in mental state)

Skin and hair

mFG score and severity
 (See Figure 2)

Ethnicity	Mild	Moderate-Severe
White/ Black	≥ 8	≥ 15
Mediterranean/ Hispanic/ Middle Eastern	≥ 9 to 10	
East Asian (e.g., China, Japan)	≥ 2 to 3	
South Asian (e.g., India, Pakistan)	≥ 8	≥ 17

Drug history

List medications and any hormonal contraceptives
 (Consider hyperandrogenic drugs e.g., testosterone, anabolic steroids, valproate)
 Access to testosterone or anabolic steroids? Yes No

Location of excess terminal hair in other areas
 (e.g., side of the face)
 Severity (0-4)

Family history

Parental consanguinity? Hirsutism CAH
 History of subfertility or infant death? PCOS T2DM
 Other.....

Presence of hypertrichosis?
 (generalised excessive hair growth in non-androgen dependent areas)

Ethnicity

Patient's ethnic origin
 How does this relate to 'expected' hair growth in patient's ethnicity?
 High risk ethnicity for CAH?
 (Ashkenazi Jew, Hispanic, Slav)

Other examination findings Acne (mild/ moderate/ severe?)
 Androgenic alopecia
 Acanthosis nigricans
 E.g., abdominal, features of thyroid dysfunction, fundoscopy, visual fields

Investigations

Previous investigations and results

Figure 1 Hirsutism assessment proforma for adolescents. BP, blood pressure; BMI, body mass index; CAH, congenital adrenal hyperplasia; d, day; DOB, date of birth; mFG, modified Ferriman-Gallwey; PCOS, polycystic ovary syndrome; QoL, quality of life; T2DM, type 2 diabetes mellitus; y, year.

Modified Ferriman-Gallwey Score Sheet

Notes: Terminal hairs are long (>5mm), coarse and dark. Excessive vellus hairs (short, fine, fair) over non-androgen dependent areas (e.g. forearms) may indicate hypertrichosis rather than hirsutism.

Recent hair removal may falsely lower Ferriman-Gallwey score. Indicate if and when hair was removed.

Name:
 DOB:
 Hospital number:

 (Or use patient label)

Body area	Degree of hair growth score	Score
Upper lip	0 No terminal hairs 1 A few terminal hairs at the outer margin 2 Small moustache at outer margin 3 Moustache extending halfway from outer margin 4 Moustache extending to midline	
Chin	0 No terminal hairs 1 A few scattered terminal hairs 2 Scattered hairs with small concentrations 3 Complete cover (light) 4 Complete cover (heavy)	
Chest	0 No terminal hairs 1 Circumareola hairs 2 Circumareola and midline hairs 3 Fusion of these areas with 3/4 cover 4 Complete cover	
Upper abdomen	0 No terminal hairs 1 A few midline terminal hairs 2 Streak of terminal hairs in the midline 3 Hair extending beyond middle, partial cover 4 Full cover	
Lower abdomen	0 No terminal hairs 1 A few midline hairs 2 Midline streak of hair 3 Midline thickened band < 1/2 width of pubic hair at base 4 Inverted V shape above pubic hair	
Upper arms	0 No terminal hairs 1 Scattered terminal hairs over less than 1/4 upper arm 2 Increased but incomplete coverage 3 Entire area covered (light growth) 4 Entire area covered (heavy growth)	
Thighs	0 No terminal hairs 1 Sparse growth covering less than 1/4 thigh 2 Sparse growth covering more than 1/4 thigh but not full coverage 3 Thigh completely covered (light) 4 Thigh completely covered (heavy)	
Upper back	0 No terminal hairs 1 Sparse terminal hairs 2 Increased numbers of spread hairs 3 Complete cover (light) 4 Complete cover (heavy)	
Lower back	0 No terminal hairs 1 Sacral tuft of terminal hair 2 Some lateral extension of hair 3 3/4 lower back covered with terminal hairs 4 Complete cover of lower back	
		Total score

Figure 2 Modified Ferriman-Gallwey score sheet.




Figure 3 Suggested adolescent hirsutism management pathway. *Testosterone values dependent on assay used and local reference ranges. 17OHP, 17-hydroxyprogesterone; COCP, combined oral contraceptive pill; CPA, cyproterone acetate; d, day; DHEAS, dehydroepiandrosterone sulfate; FAI, Free Androgen Index; mFG, modified Ferriman-Gallwey; SHBG, sex hormone-binding globulin; USP, urinary steroid profile; y, year.

testosterone levels or features of NCCAH (table 1).⁹ Patients with a moderate–severe mFG score or features of a hyperandrogenic endocrine disorder need a more detailed work-up (figure 3).

MANAGEMENT

If an underlying cause for the hirsutism has been identified, this should be treated in the first instance. Thereafter, hirsutism management falls into three categories: lifestyle, cosmetic and pharmacological. Exploring patients' expectations is important, discussing with them that no treatment will 'cure' the hirsutism and that pharmacological management can take 6–12 months to demonstrate an effect, due to the length of hair growth cycle.⁸

Lifestyle

Lifestyle interventions are particularly useful in patients with PCOS but should be discussed with all patients. If patients are overweight or obese, weight loss strategies should be discussed with signposting to appropriate support, for example, a local community weight management programme.⁸ Weight loss of 5%–7% body weight can improve testosterone levels

and menstrual irregularity, reduce the risk of metabolic sequelae and result in small improvements in hirsutism severity.^{4 8} Lifestyle changes are not as effective as pharmacological interventions and should not be used alone.⁸

Providing emotional support by psychological welfare discussions and connecting patients to peers is important. Increased social support has been correlated with higher levels of health-related quality of life in patients living with hirsutism.¹⁰ Patients can be signposted to resources such as Changing Faces (<https://www.changingfaces.org.uk>), a charity supporting people with visible differences, which offers counselling, or Verity (<http://www.verity-pcos.org.uk>), a charity run by people living with PCOS.

Cosmetic

Cosmetic measures include methods that remove hair shafts from the skin surface, depilation (eg, shaving or chemical depilation), or methods that remove hairs from above the bulb, epilation (plucking, waxing or threading). A commonly cited myth is that hair regrowth is thicker or darker after shaving. Hair may appear thicker as a blunt tip is formed rather than the tapered end of uncut hair,

but patients can be reassured that no hair removal method causes increased hair shaft diameter or darkening.⁸ Cosmetic hair removal is readily available but is temporary and can be painful, cause skin irritation, scarring, folliculitis, or hyperpigmentation, particularly in people of colour.⁸ Bleaching is a non-removal alternative generally used on facial hair.

Topical

Eflornithine cream inhibits hair growth in the anagen phase (growing stage) of the hair cycle and is licensed for facial use.¹¹ Noticeable effects take 6–8 weeks, but hair growth returns to pretreatment levels when stopped, and it can cause itching and dry skin.⁸

Longer-lasting hair reduction

Photoepilation and electrolysis are methods of permanently reducing hair counts. Photoepilation, or 'laser' treatment, uses light pulses to thermolyse pigmented terminal hair follicles and can reduce hair by 80% after 6 months of treatment.¹² It should be avoided in patients from Mediterranean or Middle Eastern heritage because of potential paradoxical hypertrichosis.⁸ Photoepilation is up to 60 times faster than electrolysis, as electrolysis uses a fine electrode to thermolyse each individual hair. It is more painful, limited to smaller areas and not available on the NHS.¹² Funding approval for NHS laser therapy is very limited. Private clinics may offer photoepilation but, in our experience, will not accept patients <18 years. Less powerful photoepilation devices are available to purchase for home use but are expensive and not as effective as medical photoepilation devices.⁸ Eflornithine cream can be an effective adjunct, facilitating a more rapid response.⁸

Pharmacological

Combined oral contraceptive pills (COCPs)

COCPs are first-line treatment for hirsutism, although they are prescribed off-label. They slow hair growth but do not remove existing hair.⁸ The oestrogen decreases free androgens by increasing SHBG production and suppresses luteinising hormone, reducing adrenal and ovarian androgen production. The progestins block androgen receptors and conversion of testosterone to the more potent dihydrotestosterone by inhibition of 5 α -reductase.⁴ There is no clear evidence for choosing any particular COCP; however, third-generation progestins, for example, desogestrel, gestodene and norgestimate, are 'androgen neutral' and are preferred to second-generation progestins like levonorgestrel. Antiandrogenic-containing COCPs, including cyproterone acetate or drospirenone, improve hirsutism more than other COCPs.^{8,13}

Ethinylestradiol-containing COCPs are preferred to 17- β -estradiol or estradiol valerate, which are less likely to suppress ovarian androgens due to their lower oestrogen dose.⁸ Patients should be made aware of the increased risk of thromboembolism with COCPs and the related symptoms but reassured that the absolute risk is very small.¹³

Test your knowledge

- Which of the following are features of non-classic congenital adrenal hyperplasia? You may choose more than one answer.
 - Increased muscle mass.
 - Advanced bone age.
 - Hyponatraemia.
 - Raised 21-hydroxylase.
 - Raised 17-hydroxyprogesterone.
- A 15-year-old patient attends with concerns about dark hairs on her chin and abdomen. You find mild hirsutism on examination. She attained menarche at 11 years old and thinks she has only had two periods in the last year. You suspect she has polycystic ovary syndrome (PCOS). Which of the following is the most appropriate approach to investigation?
 - Pelvic ultrasound.
 - No investigations needed, PCOS is a clinical diagnosis.
 - Testosterone and sex hormone-binding globulin.
 - Androgen panel (17OHP, androstenedione, DHEAS and testosterone).
 - No investigations needed, unable to diagnose PCOS at this age.
- Regarding pharmacological treatment of hirsutism, which statement is the most correct?
 - Flutamide is a second-line treatment.
 - Oral contraceptive use alone will get decrease existing terminal hair.
 - Metformin is a useful adjunct for the treatment of hirsutism in PCOS.
 - The antiandrogenic effect of spironolactone is dose dependent.
 - The progesterone only contraceptive pill can be used if the combined pill is contraindicated or not tolerated.
- Which of following statements about cosmetic treatments of hirsutism is correct?
 - Eflornithine cream works by disrupting the telogen, or resting, phase of hair growth.
 - Photoepilation is much faster than electrolysis.
 - Shaving causes hair to grow back thicker.
 - Photoepilation is effective on pale coloured hairs.
 - COCPs should be used for at least 9 months before another treatment is considered.
- Which of the following may be features of an androgen secreting tumour? You may choose more than one answer.
 - Insidious onset of symptoms.
 - Voice change.
 - Suppressed DHEAS.
 - Raised androstenedione.
 - Progression of hirsutism despite treatment.

Answers to the quiz are at the end of the references.

Antiandrogens

Antiandrogens are second-line treatments if COCPs are contraindicated, not tolerated or ineffective (ie, 6 months without improvement).^{4,8} They may be used in conjunction with COCPs as part of first-line therapy in moderate to severe hirsutism.⁸ Spironolactone and finasteride are the recommended antiandrogens in adolescents but should be initiated under specialist guidance.^{3,8} Spironolactone inhibits the androgen receptor and 5 α -reductase.⁸ Its antiandrogenic effects are dose dependent, and the recommended starting dose is 25 mg.⁸ Electrolytes should be monitored due to risk of hyperkalaemia. It should not be used with drospirenone-containing COCPs due to the cumulative antiminerocorticoid effect.⁸ Finasteride is a partial 5 α -reductase inhibitor. Antiandrogens can cause feminisation of a male foetus and should be prescribed with effective contraception and appropriate counselling.¹⁴ Flutamide is an effective antiandrogen but can be hepatotoxic and not recommended.⁸

Metformin is an insulin sensitiser used in PCOS for its metabolic effects and improvement in menstrual regularity but has negligible effect on hirsutism.⁸

CONCLUSION

Hirsutism commonly affects adolescent girls and necessitates careful assessment and tailored investigation. Management varies, depending on the severity, aetiology and individual patient factors. The psychological and social impact of hirsutism should be sought in all and appropriate support offered. Working with patients to understand the chronic nature of hirsutism and likely outcome of the available therapies may improve patient satisfaction.

Twitter Charlotte Jane Elder @charlottejelder

Contributors ASC designed, drafted and revised all aspects of the work, approved the final version for publishing and agreed to be accountable for all aspects of the work. CJE conceptualised, designed and revised all aspects of the work, approved the final version for publishing and agreed to be accountable for all aspects of the study. MM conceptualised and revised the work, particularly regarding cosmetic management of hirsutism, approved the final version for publishing and agreed to be accountable for all aspects of the study.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Commissioned; externally peer reviewed.

ORCID iDs

Alexandra Selina Cross <http://orcid.org/0000-0002-9799-582X>

Charlotte Jane Elder <http://orcid.org/0000-0003-2390-5593>

REFERENCES

- 1 Yildiz BO. Assessment, diagnosis and treatment of a patient with hirsutism. *Nat Clin Pract Endocrinol Metab* 2008;4:294–300.
- 2 Rosenfield RL. The diagnosis of polycystic ovary syndrome in adolescents. *Pediatrics* 2015;136:1154–65.
- 3 Esquivel-Zuniga MR, Kirschner CK, McCartney CR, *et al.* Non-PCOS Hyperandrogenic disorders in adolescents. *Semin Reprod Med* 2022;40:42–52.
- 4 Witchel SF, Oberfield SE, Peña AS. Polycystic ovary syndrome: pathophysiology, presentation, and treatment with emphasis on adolescent girls. *J Endocr Soc* 2019;3:1545–73.
- 5 Ibáñez L, Oberfield SE, Witchel S, *et al.* An international consortium update: pathophysiology, diagnosis, and treatment of polycystic ovarian syndrome in adolescence. *Horm Res Paediatr* 2017;88:371–95.
- 6 Hatch R, Rosenfield RL, Kim MH, *et al.* Hirsutism: implications, etiology, and management. *Am J Obstet Gynecol* 1981;140:815–30.
- 7 Legro RS, Arslanian SA, Ehrmann DA, *et al.* Diagnosis and treatment of polycystic ovary syndrome: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2013;98:4565–92.
- 8 Martin KA, Anderson RR, Chang RJ, *et al.* Evaluation and treatment of hirsutism in premenopausal women: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2018;103:1233–57.
- 9 Carmina E, Dewailly D, Escobar-Morreale HF, *et al.* Non-classic congenital adrenal hyperplasia due to 21-hydroxylase deficiency revisited: an update with a special focus on adolescent and adult women. *Hum Reprod Update* 2017;23:580–99.
- 10 Ekbäck MP, Lindberg M, Benzein E, *et al.* Social support: an important factor for quality of life in women with hirsutism. *Health Qual Life Outcomes* 2014;12:183.
- 11 Bode D, Seehusen DA, Baird D. Hirsutism in women. *Am Fam Physician* 2012;85:373–80.
- 12 Kutlubay Z. Alexandrite laser hair removal results in 2359 patients: a Turkish experience. *J Cosmet Laser Ther* 2009;11:85–93.
- 13 Escobar-Morreale HF, Carmina E, Dewailly D, *et al.* Epidemiology, diagnosis and management of hirsutism: a consensus statement by the androgen excess and polycystic ovary syndrome Society. *Hum Reprod Update* 2012;18:146–70.
- 14 Teede HJ, Misso ML, Costello MF, *et al.* Recommendations from the International evidence-based guideline for the assessment and management of polycystic ovary syndrome. *Hum Reprod* 2018;33:1602–18.
- 15 Carmina E, Rosato F, Janni A, *et al.* Extensive clinical experience: relative prevalence of different androgen excess disorders in 950 women referred because of clinical hyperandrogenism. *J Clin Endocrinol Metab* 2006;91:2–6.
- 16 Fleseriu M, Auchus R, Bancos I, *et al.* Consensus on diagnosis and management of Cushing's disease: a guideline update. *Lancet Diabetes Endocrinol* 2021;9:847–75.
- 17 Fahie-Wilson M, Smith TP. Determination of prolactin: the macroprolactin problem. *Best Pract Res Clin Endocrinol Metab* 2013;27:725–42.
- 18 Sharma D, Shanker V, Tegta G, *et al.* Clinico-investigative profile of patients of hirsutism in a tertiary level institution. *Int J Trichology* 2012;4:69.

Answers to the multiple choice questions

1. (A) True; (B) True; (C) False; (D) False; (E) True.
2. (A) False; (B) False; (C) True; (D) False; (E) False.
3. (A) False; (B) False; (C) False; (D) True; (E) False.
4. (A) False; (B) True; (C) False; (D) False; (E) False.
5. (A) False; (B) True; (C) False; (D), True; (E) True.