



OPEN ACCESS

Open Access
Scan to access more
free content

Fifteen-minute consultation: an evidence-based approach to research without prior consent (deferred consent) in neonatal and paediatric critical care trials

Kerry Woolfall,¹ Lucy Frith,² Angus Dawson,³ Carrol Gamble,⁴ Mark D Lyttle,^{5,6} the CONNECT advisory group Bridget Young¹

¹Department of Psychological Sciences, University of Liverpool, Liverpool, UK

²Department of Health Service Research, University of Liverpool, Liverpool, UK

³Centre for Values, Ethics and the Law in Medicine (VELiM), School of Public Health, University of Sydney, Sydney, New South Wales, Australia

⁴Department of Biostatistics, University of Liverpool, Liverpool, UK

⁵Emergency Department, Bristol Royal Hospital for Children, Bristol, UK

⁶Faculty of Health and Applied Sciences, University of the West of England, Bristol, UK

Correspondence to

Dr Kerry Woolfall, Department of Psychological Sciences, University of Liverpool, Liverpool L69 3GL, UK; K.Woolfall@liverpool.ac.uk

Received 26 June 2015

Revised 11 September 2015

Accepted 12 September 2015

Published Online First

13 October 2015

WHAT DO WE MEAN BY RESEARCH WITHOUT PRIOR CONSENT (DEFERRED CONSENT)?

Emergency research with critically unwell children is vital to make sure that the most ill and injured children benefit from evidence-based healthcare.¹ Ethical guidance require that consent be sought from parents (or legal representatives) on behalf of their children² before research is initiated, yet concerns about problems in seeking parents' consent when their child is critically ill have been a significant barrier to conducting clinical trials.³⁻⁴ Taking time out to seek informed consent before starting treatment will often be difficult to justify as delaying any intervention in an emergency could diminish a child's chances of recovery. Parents will usually be highly distressed in a critical care situation, and many will struggle to make an informed decision about research in the limited time available.

Many countries have legislated to permit variations to informed consent and allow progress in research to develop critical care treatments.⁵⁻⁷ While the details vary, a common feature is that informed consent is not requested before the patient receives the intervention being researched.⁸ In the USA, the Food and Drug Administration (FDA) Exception from Informed Consent (EFIC) essentially 'waives' informed consent, although practitioners must show that they have attempted to contact legal representatives and tried to provide the opportunity to 'opt out' of a trial.⁵⁻⁹ The FDA's detailed guidance aims to assist researchers in implementing

EFIC,¹⁰⁻¹¹ although the accompanying public consultation requirements have led to varied practice and costly delays in setting up trials.¹²

Across European Union countries, legislation⁷⁻¹³ enables practitioners to conduct research without seeking prior informed consent from parents when certain conditions are met (see [box 1](#) for UK example). No accompanying guidance has been made available to assist researchers in implementing the legislation. European legislation does not name this alternative to informed consent, but it is commonly called 'deferred consent'. We would argue that this is a misnomer as a child will have already received an intervention as part of a trial before any information is given or consent is sought. Essentially permission is sought post-intervention to use data that have already been collected and consent for the child to continue to take part in the trial. These problems with the terminology have led to much discussion recently, leading to a move towards the term 'research without prior consent', as it more accurately reflects the process of consent seeking in critical care research. We will, therefore, use the latter term for the rest of this article. However, regardless of what terminology is used, research without prior informed consent can be seen as eroding the autonomy of parents and children and has been much debated.¹⁴⁻¹⁸

HAVE ANY TRIALS BEEN CONDUCTED WITHOUT PRIOR CONSENT?

Although a number of adult critical care trials have been conducted without prior



CrossMark

To cite: Woolfall K, Frith L, Dawson A, et al. *Arch Dis Child Educ Pract Ed* 2016;**101**:49–53.

Box 1 Research without prior consent (deferred consent) can be conducted when the following conditions are met

1. Treatment is required urgently
2. Urgent action is required for the purposes of the trial
3. It is not reasonably practicable to obtain consent prospectively
4. An ethics committee has given approval to the procedure under which the action is taken.¹³

consent over the last decade,^{19 20} the CAThether infections in CHildren trial (CATCH)²¹ was one of the first UK paediatric trials to use this approach since legislation changes in 2008. Another National Institute for Health Research-funded multicentre trial investigating the Emergency treatment with Levetiracetam or Phenytoin in Status Epilepticus,^{22 23} which is using research without prior consent, has recently opened to recruitment. The number of trials using this approach is expected to increase over the next few years. It will be important to share experiences of conducting these challenging trials to inform peer and Research Ethics Committee reviews.

WHAT DO PARENTS AND PRACTITIONERS THINK ABOUT RESEARCH WITHOUT PRIOR CONSENT?

In the CONsent methods in childreN's emergEncy medicine and urgent Care Trials (CONNECT) study, we found that many parents recruited to CATCH were momentarily shocked or surprised to discover that their child had been entered into a trial without their consent, although they did not voice this to practitioners.²⁴ After hearing practitioners explain why research without prior consent is being used in their situation—that it enables vital research to take place in time-critical situations—parents' initial concerns were dispelled. Practitioners' explanations were important to parents and helped to reassure them that there were good reasons for doing research without prior consent. Gauging the right moment to approach parents to discuss research is important—we found that mistiming the approach could add to parents' distress.

Despite its 'do then ask' sequence, parents with experience of research without prior consent told us they felt *their* decisions about their child's participation had been voluntary. CONNECT and other research^{25 26} have shown that parents support research without prior consent and appreciate the reasons for using it as long as their child's safety is not compromised. However, parents' support for this approach may have its limits and is related to what is being trialled. Most parents in CONNECT remarked that they would be concerned about not seeking prior consent in trials involving either 'new' drug

interventions that were not already used in clinical care or other potentially significant changes in clinical practice.^{23 24}

Practitioners' views on research without prior consent differed depending upon whether or not they had experience of this method.²⁷ Practitioners with no experience of research without prior consent were concerned that it would be detrimental to the parent–practitioner relationship. In contrast, practitioners with experience of this approach described how families were receptive to the method as long as discussions were appropriately timed and conducted sensitively.

We drew on the CONNECT study findings in light of bioethical principles, including voluntariness, autonomy, non-maleficence and justice,^{18 28} to produce guidance on approaches to critical care research without prior informed consent.

WHEN SHOULD I APPROACH PARENTS TO DISCUSS RESEARCH WITHOUT PRIOR CONSENT?

CONNECT guidance recommends explaining what has happened at the earliest *appropriate* opportunity, which is likely to be after the initial emergency situation has passed. In such clinical situations, parents often rapidly form a close relationship with the child's nurse. Consulting with nursing staff about the child's

Box 2 Key points to cover when discussing research without prior consent with parents

- ▶ Why the research is being conducted and why their child's condition made him/her eligible for the trial.
- ▶ That it was not possible to seek consent before the research intervention was given because their child needed immediate treatment, and it was not safe to delay this.
- ▶ That their permission is being sought to use information and/or samples that have already been collected and for their child to continue in the research.
- ▶ Details of how the intervention is already used in clinical practice (if applicable), any changes to clinical practice and potential risks of being in the research.
- ▶ That the research has been approved by an independent research ethics committee whose role is to review research to protect the rights, safety and well-being of participants.
- ▶ How the research findings will inform future treatments for critically ill children.
- ▶ That parents are free to choose whether or not their child's information is used in the research and that their decision will not affect their child's care.
- ▶ Details of any follow-up procedures arising from the research (if applicable).
- ▶ Where further information can be found—for example, leaflets, website, principal investigator.

condition and how parents are coping will help gauge when is an appropriate time.

HOW SHOULD I EXPLAIN TO PARENTS THAT THEIR CHILD HAS BEEN ENTERED INTO RESEARCH WITHOUT THEIR PRIOR INFORMED CONSENT?

Ask a member of staff known to the family to introduce you. Start by asking parents how their child is doing and check whether it is a convenient time to discuss research. Discuss key aspects of the trial, showing parents (and children if appropriate—see below) where information can be found on the

participant information leaflet, paying particular attention to the key points shown in [box 2](#). Allow parents time to consider the information (overnight if possible) and to ask questions about the trial. While it may be important to seek permission to use data already collected and consent for continued participation in the trial (and any follow-up procedures) before the child is discharged from hospital, it is also important to allow time to explore parents' views and understanding of the trial and follow-up procedures.

Be prepared to address concerns that participation may have contributed to a poor recovery. It may help to explain any potential risks associated with

Box 3 Options to consider when a child has died

Option 1: Approach parents to explain about the trial before they leave hospital

- ▶ Discuss the trial and provide information before parents leave hospital. However, only approach parents with information and seek permission to use data already collected at this point if it is believed that parents have the capacity to absorb information and make an informed decision.

Option 2: Explain about the trial by letter at a later date

- ▶ If it is not thought appropriate to explain about the trial or seek permission to use data already collected before parents leave the hospital, consult with clinical colleagues and bereavement counsellors to identify an appropriate time to contact parents via a posted letter. Sending the letter could be timed to coincide with the bereavement follow-up invitation.
- ▶ The covering letter, information leaflet and consent form should be designed and worded specifically for bereaved parents. It should be prepared at the trial design stage and written in close consultation with bereaved parents, bereavement specialists and relevant special interest groups (see recommendation 1).
- ▶ The covering letter should be personalised and, if possible, signed by a clinician known to the family. The letter should explain that, understandably, parents will often have questions about the research in the days, weeks or months after the loss of a child and invite them to contact the trial team to arrange for a telephone or face-to-face discussion with the principal investigator about the trial if they wish. Include the bereaved parent information leaflet, consent form and stamped addressed envelope.
- ▶ At the outset of the trial, ethical approval may have been sought to include the anonymised data of deceased patients in analyses should no consent form be received from bereaved parents. Therefore, letters to parents should explain whether or not their child's data will be included in the trial if parents do not respond to the letter.
- ▶ Copies of the letters and accompanying documents sent to parents should be placed in the patient's notes.
- ▶ Be prepared to respond to parents who are concerned that research participation may have contributed to their child's death. Be careful to avoid giving false reassurance that this is not the case, unless it has been established by the principal/chief investigator that the cause of death was not related to the trial.

Option 3: Contact parents by telephone or letter to arrange a face-to-face discussion about the trial

- ▶ If it is not thought appropriate to explain about the trial or seek permission to use data already collected before parents leave the hospital, consult with clinical colleagues and bereavement counsellors to identify an appropriate time to contact parents via telephone or letter to arrange a face-to-face visit to discuss research.
- ▶ The letter should be personalised, signed by a clinician (known to the family if possible) and include a bereaved parent information leaflet.
- ▶ Copies of letters sent to parents should be placed in the patients' notes.
- ▶ Provide parents with options for meeting location (eg, at their home or local hospital) as some parents may not wish to return to the hospital where their child died.
- ▶ During face-to-face discussions, explore parents' views and understanding of the trial and why consent was not sought so that any concerns can be addressed.
- ▶ Be prepared to respond to parents who are concerned that trial participation may have contributed to their child's death. Be careful to avoid giving false reassurance that the trial did not contribute to their child's death unless it has been established by the principal/chief investigator that the cause of death was not related to the trial.
- ▶ If parents do not wish to have a face-to-face meeting, inform them that a trial information leaflet and consent form will be sent via post (see option 2).

participation in the trial, that the intervention is already used in clinical practice (if applicable), and to indicate that the research has been approved by a Research Ethics Committee. It may also help to explain that nobody will know which treatment is the most effective until the trial has been completed (which may take a few years) and to offer parents the opportunity to speak to the principal investigator or senior member of the research team to discuss any concerns.

SHOULD I INVOLVE CHILDREN IN THE DISCUSSION?

Although children (under 16 years) cannot legally provide consent for their own participation in a trial, decisions about research should be shared by children and their parents²⁹ if their maturity, condition and cognitive capacity allows. Young people (aged 16–18 years)³⁰ can legally provide their own consent for a trial, although this is often impossible in an emergency situation. When assent (for children) or consent (for young people) cannot be sought due to their clinical condition, provide a developmentally appropriate information sheet to help parents discuss the research with their child when they have recovered. Provide contact details so that parents or children can discuss any aspect of the trial with the research team at a later date if they wish.

WHAT SHOULD HAPPEN IF A CHILD DIES BEFORE PARENTS ARE APPROACHED ABOUT THE TRIAL?

Legislation^{7 13} does not stipulate what should happen in circumstances where a child dies. As there is wide variability and complexity of parental feelings about research when a child has been enrolled in a study and subsequently died, a one-size-fits-all approach to discussing a clinical trial is unlikely to be sensitive to the needs of grieving parents.³¹ Although there are some exceptions,²⁵ many bereaved parents wish to be informed about their child's involvement and provided with the opportunity to discuss having their child's data analysed in a trial.^{23 31} Talking with recently bereaved parents about research in which their child was involved will almost certainly be very difficult for you and for parents. However, in the interests of openness and honesty it is important to offer the opportunity to do so, otherwise parents will have no knowledge of their child's participation in research, nor can their child's data be included in the analyses. As well as potentially biasing the findings,³² this could be contrary to what parents would want for their child's data. **Box 3** outlines some options to help those involved decide how to approach bereaved parents to discuss research without prior consent.

CONCLUSIONS

The CONNECT guidance will help practitioners to conduct research without prior consent in a way that is

ethically appropriate and addresses the needs of families. Full CONNECT guidance can be found at <https://www.liv.ac.uk/psychology-health-and-society/research/connect/> and will be reviewed and updated as further evidence becomes available. Research is required to look at the transferability of CONNECT guidance to other study types and settings, including adult critical care.

Twitter Follow Kerry Woolfall at @CONNECTStudy, Follow Mark Lyttle at @mdlyttle

Acknowledgements We thank all the parents and practitioners who shared their experiences with us. We are grateful to the CATCH trial team and all the CATCH doctors and nurses for their help with recruitment to CONNECT.

Collaborators Julia Harris, Michael Parker, Claire Snowdon, Helen Hickey and Hazel Greig-Midlane.

Contributors KW, LF, AD, CG, MDL and BY wrote the manuscript. All authors, including the CONNECT advisory group, developed the CONNECT guidance, critically appraised and approved the final manuscript.

Funding CATCH was funded by the National Institute for Health Research Health Technology Assessment (NIHR HTA) programme (project number 08/13/47). CONNECT was funded by Wellcome Trust (WT095874MF) and supported by the MRC Network of Hubs for Trials Methodology Research (MR/L004933/1- R/N42).

Competing interests None declared.

Ethics approval Northwest Liverpool East Research Ethics Committee: 12/NW/0094.

Provenance and peer review Commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Kanthimathinathan HK, Scholefield BR. Dilemmas in undertaking research in paediatric intensive care. *Arch Dis Child* 2014;99:1043–9.
- 2 Helsinki WEU-Do. World medical association declaration of Helsinki. In: Assembly WG, ed. Fortaleza, Brazil, 2013. <http://www.wma.net/en/30publications/10policies/b3/> (accessed 10 Sept 2015).
- 3 Roberts I, Prieto-Merino D, Shakur H, *et al.* Effect of consent rituals on mortality in emergency care research. *Lancet* 2011;377:1071–2.
- 4 Maitland K, Molyneux S, Boga Mwamvua, *et al.* Use of deferred consent for severely ill children in a multi-centre phase III trial. *Trials* 2011;12:90.
- 5 US Department of Health and Human Services Food and Drug Administration. *Guidance for Institutional Review Boards, Clinical Investigators, and Sponsors. Exception from informed consent requirements for emergency research.* Rockville, MD: US Department of Health and Human Services Food and Drug Administration, 2013.
- 6 National Health and Medical Research Council. *National Statement on Ethical Conduct in Human Research (2007): updated May 2015.* Australia: National Health and Medical Research Council, 2007.
- 7 The European Parliament and the Council of the European Union. Regulation (EU) No 536/2014 of the European Parliament and the council of 16 April 2014 on clinical trials

- on medicinal products for human use, and repealing Directive 2001/20/EC: Official Journal of the European Union, 2014.
- 8 Jansen-van der Weide M, Caldwell P, Young B, *et al.* Clinical trial decisions in difficult circumstances: parental consent under time pressure. *Pediatrics* 2015;eFirst. doi:10.1542/peds.2014-3402
 - 9 Eltorki M, Uleryk E, Freedman SB. Waiver of informed consent in pediatric resuscitation research: a systematic review. *Acad Emerg Med* 2013;20:822–34.
 - 10 Shamoo AE. Letter to the editor: emergency research consent waiver—a proper way. *Am J Bioeth* 2006;6:W48–51.
 - 11 McClure KB, DeIorio NM, Gunnels MD, *et al.* Attitudes of Emergency Department Patients and Visitors Regarding Emergency Exception from Informed Consent in Resuscitation Research, Community Consultation, and Public Notification. *Acad Emerg Med* 2003;10:352–9.
 - 12 Richardson LD, Quest TE, Birnbaum S. Communicating with communities about emergency research. *Acad Emerg Med* 2005;12:1064–70.
 - 13 Legislation.gov.uk. The Medicines for Human Use (Clinical Trials) and Blood Safety and Quality (Amendment) Regulations 2008 941. 10. 2008.
 - 14 Cooke RWI. Good practice in consent. *Semin Fetal Neonatal Med* 2005;10:63–71.
 - 15 Kottow M. The battering of informed consent. *J Med Ethics* 2004;30:565–9.
 - 16 Brierley J, Larcher V. Emergency research in children: options for ethical recruitment. *J Med Ethics* 2011;37:429–32.
 - 17 Nelson RM, Beauchamp T, Miller VA, *et al.* The concept of voluntary consent. *Am J Bioeth* 2011;11:6–16.
 - 18 Beauchamp T, Childress J. *Principles of biomedical ethics*. Oxford: Oxford University Press, 2001.
 - 19 Perkins GD, Lall R, Quinn T, *et al.* Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomised controlled trial. *Lancet* 2015;385:947–55.
 - 20 Edwards P, Arango M, Balica L, *et al.*, Crash trial collaborators. Final results of MRC CRASH, a randomised placebo-controlled trial of intravenous corticosteroid in adults with head injury—Outcomes at 6 months. *Lancet* 2005;365:1957–9.
 - 21 CTRC. *Catheter Infections in Children (CATCH) Protocol V4.0*. Liverpool, 2011. <http://www.catchtrial.org.uk/publications.html>
 - 22 National Institute for Health Research. HTA—12/127/134: A pragmatic randomised controlled trial of intravenous levetiracetam versus intravenous phenytoin in terminating acute, prolonged tonic clonic seizures including convulsive status epilepticus in children, the ECLIPSE Study: Emergency treatment with Levetiracetam or Phenytoin in Status Epilepticus. *EvalTrials Stud* 2015. <http://www.nets.nihr.ac.uk/projects/hta/12127134> (accessed 24 Sep 15).
 - 23 Woolfall K, Young B, Frith L, *et al.* Doing challenging research studies in a patient-centred way: a qualitative study to inform a randomised controlled trial in the paediatric emergency care setting. *BMJ Open* 2014;4:e005045.
 - 24 Woolfall K, Frith L, Gamble C, The CONNECT advisory group, *et al.* How parents and practitioners experience deferred consent for emergency research: a mixed method study. *BMJ Open* 2015;5:e008522. doi:10.1136/bmjopen-2015-008522
 - 25 Gamble C, Nadel S, Snape D, *et al.* What parents of children who have received emergency care think about deferring consent in randomised trials of emergency treatments: postal survey. *PLoS ONE* 2012;7:e35982.
 - 26 Molyneux S, Njue M, Boga M, *et al.* ‘The words will pass with the blowing wind’: staff and parent views of the deferred consent process, with prior assent, used in an emergency fluids trial in two African Hospitals. *PLoS ONE* 2013;8:e54894.
 - 27 Woolfall K, Frith L, Gamble C, *et al.* How experience makes a difference: Practitioners’ views on the use of deferred consent in paediatric and neonatal emergency care trials. *BMC Med Ethics* 2013;14:45.
 - 28 Frith L. Symbiotic Empirical Ethics: A Practical Methodology. *Bioethics* 2012;26:198–206.
 - 29 Nuffield Council on Bioethics. *Children and clinical research: ethical issues*. London: Nuffield Council on Bioethics, 2015.
 - 30 Legislation.gov.uk. The Medicines for Human Use (Clinical Trials) Amendment (No. 2) Regulations (2006) 10. 2006.
 - 31 Snowdon C, Brocklehurst P, Tasker R, *et al.* Death, Bereavement and randomised controlled trials (BRACELET): a methodological study of policy and practice in neonatal and paediatric intensive care trials. *Health Technol Assess* 2014;18:1–410.
 - 32 Jansen TC, Kompanje EJ, Druml C, *et al.* Deferred consent in emergency intensive care research: what if the patient dies early? Use the data or not? *Intensive Care Med* 2007;33:894–900.